

DAIDS Grading for Adverse Events



NAUSEA & VOMITING

- mild nausea & vomiting
- without any symptoms of dehydration

- slightly more nausea & vomiting
- with symptoms of mild dehydration (increased thirst, dry tongue, mouth, etc.) and
- generalized weakness

- excessive vomiting,
- weakness and inability to stand,
- sunken eyes with loss of skin elasticity and
- electrolyte imbalance

- all symptoms of 'Severe Category',
- blood in vomit with
- altered level of consciousness



GASTRITIS & PAIN IN ABDOMEN

- mild pain in the upper abdominal region, discomfort

- acid reflux, sour taste in mouth
- burning sensation in upper abdominal region

- severe pain in the abdomen
- intensified acidity symptoms
- vomiting

- symptoms of severe acidity with or without blood in vomit & stool
- associated with altered level of consciousness
- with electrolyte imbalance



DIARRHOEA

- 2-3 loose stools
- no symptoms of dehydration

- 3-10 loose stools or watery stools with
- symptoms of mild to moderate dehydration

- >10 watery stools with
- symptoms of severe dehydration & electrolyte imbalance

- watery stools with symptoms of severe dehydration and
- blood in stool, fever, intense abdominal pain
- may or may not be associated with altered level of consciousness



PERIPHERAL NEUROPATHY

- mild numbness & weakness in hands & feet

- prickling, stabbing, burning or tingling sensation along with
- gradual increase in numbness & weakness

- symptoms of moderate neuropathy along with
- extreme sensitivity to touch

- muscle weakness with lack of coordination and difficulty to balance,
- there may be poor control of bowel & bladder movement



ARTHRALGIA (JOINT PAIN)

MILD	MODERATE	SEVERE	POTENTIALLY LIFE-THREATENING
<ul style="list-style-type: none"> • Pain on touching joint • none or minimal interference with usual social & functional activities 	<ul style="list-style-type: none"> • pain while walking • swelling, redness, warmth in and around the involved joint • causing greater than minimal interference with usual social & functional activities 	<ul style="list-style-type: none"> • stiffness of the involved joint, along with • signs of increased tenderness • causing inability to perform usual social & functional activities 	<ul style="list-style-type: none"> • generalized weakness and limited movement due to significant stiffness & tenderness of the joint • causing inability to perform basic self-care functions



TENDONITIS (ACHILLES TENDON)

<ul style="list-style-type: none"> • pain, swelling & stiffness along the Achilles tendon, in the morning 	<ul style="list-style-type: none"> • pain along the tendon or back of the heel that worsens with movement 	<ul style="list-style-type: none"> • severe pain along the tendon/back of the heel leading to inability to walk 	<ul style="list-style-type: none"> • severe pain along the tendon/back of the heel, weakening of tendon which may lead to rupture
--	--	--	--



SKIN RASHES, ALLERGIC REACTIONS & ANAPHYLAXIS

<ul style="list-style-type: none"> • mild urticaria (itching and skin rashes) but localised to smaller part of the body 	<ul style="list-style-type: none"> • generalized itching or covering a significant area of the body • skin rashes with tingling & burning sensations 	<ul style="list-style-type: none"> • intolerable itching, • skin rashes in the form of raised weals of bigger size, • may be angioedema with swelling of lips & tongue 	<ul style="list-style-type: none"> • serious allergic reaction in the form of Anaphylaxis*/ Stevens–Johnson syndrome**/Lyell's syndrome***
--	--	---	---




HYPOTHYROIDISM

<ul style="list-style-type: none"> • loss of energy • dry & itchy skin • mild constipation 	<ul style="list-style-type: none"> • fatigue and generalized weakness, • unexplained weight gain, • intolerance to cold, • constipation, • swelling in the front of neck 	<ul style="list-style-type: none"> • all symptoms of hypothyroidism in severe form, • loss of appetite • drowsiness • patient may go into a state of depression 	<ul style="list-style-type: none"> • palpitation with low heart rate, increased blood pressure, • generalized edema with puffiness of face, patient may go into coma
---	---	---	--



TINNITUS AND DIZZINESS

MILD	MODERATE	SEVERE	POTENTIALLY LIFE-THREATENING
<ul style="list-style-type: none"> intermittent ringing & buzzing in ears (unilateral/bilateral), fullness in ears 	<ul style="list-style-type: none"> continuous ringing in ears patient feels nauseated & may vomit. loss of concentration due to drowsiness and giddiness 	<ul style="list-style-type: none"> symptoms of moderate tinnitus difficulty in maintaining balance 	<ul style="list-style-type: none"> increased severity of symptoms with inability to sleep
 <h2>PSYCHIATRIC BEHAVIOUR</h2> <ul style="list-style-type: none"> social withdrawal and isolation due to TB and its associated taboos a sense of hopelessness lack of self-care depression none or minimal interference with usual social and functional activities 	<ul style="list-style-type: none"> depression, disorientation, unable to concentrate causing greater than minimal interference with usual social & functional activities 	<ul style="list-style-type: none"> hallucination, delusion, disorientation in thought, speech & behaviour; severe depression causing inability to perform usual social & functional activities 	<ul style="list-style-type: none"> all symptoms of psychotic behaviour in more severe form causing inability to perform basic self-care functions sometimes, severe depression may lead to suicidal attempt